

## Functions of form and points of integration finite elements

---

### Summary:

One describes the geometry and the topology of the finite elements established in *Code\_Aster* ; for each element of reference, the expression of the functions of form and the various families of points of integration as well as the associated weights are detailed.

## Contents

---

1 Introduction.....	3
2 Linear elements: SE2, SE3 and SE4.....	4
3 Surface elements.....	6
3.1 Triangles: ELREFE TR3, TR6, TR7.....	6
3.2 Quadrangles: ELREFE QU4, QU8, QU9.....	10
4 Voluminal elements.....	13
4.1 Tetrahedrons: ELREFE TE4, T10.....	13
4.2 Pentahedrons: ELREFE PE6, P15, P18.....	15
4.3 Hexahedrons: ELREFE HE8 , H20 , H27.....	19
4.4 Pyramids: ELREFE PY5, P13.....	24
5 Bibliography.....	28
6 History of the versions of the document.....	28

## 1 Introduction

In *Code\_Aster*, one calls "finite element", a triplet (phenomenon, modeling, type of mesh). There are three principal phenomena: MECHANICS, THERMICS and ACOUSTICS.

There exist many modelings; for example, for the phenomenon MECHANICS : 3D, C\_PLAN, D\_PLAN, AXIS, DKT, POU\_D\_E,...

For a given modeling (for example 3D) of a phenomenon (for example MECHANICS), there in general exist several finite elements: an element by type of mesh supported: HEXA8, HEXA20, PENTA6,...

With final, there thus exists of very many finite elements (more than 500 in July 2004).

On the other hand, the types of mesh are them of reduced number: POI1, SEG2, SEG3, SEG4, TRIA3, TRIA6, TRIA7, QUAD4, QUAD8, HEXA8, HEXA20, ..., TETRA4, TETRA10.

In general, each finite element, to carry out its elementary calculations, uses the concepts of function of interpolation (or function of form) and of diagram of integration. In general also, these functions of form and these diagrams of integration are defined on an element known as "of reference" whose geometry is defined in an often called frame of reference:  $(\xi, \eta, \zeta)$ . The passage of the element of reference to the real element is done thanks to a geometrical transformation which uses the same functions of interpolation. The element is then known as "isoparametric". These concepts are very well explained in [bib1].

The high number of finite elements of the code combined with the restricted number of the types of mesh, conduit to the fact that there exist several finite elements being based on the same type of mesh; for example the quadrilateral with 8 nodes called QUAD8 support more than 60 different finite elements.

In theory, each finite element can choose its functions of interpolation and its diagrams of integration as he hears it. But in practice, almost all the finite elements being based on the same type of mesh, use the same element of reference, the same functions of form and the same diagrams of integration. The goal of this document is to describe these various elements of reference.

For each element of reference (called in the continuation of the document ELREFE), one will indicate:

- the mesh support, the number of the nodes, their local classification and their coordinates,
- algebraical expressions of the functions of form and their derivative first (and sometimes seconds)
- families of points of integration which one will name. For each family, one will give the number of points, their coordinates and their "weights" of integration. The sum of these weights, must give the "volume" of the element of reference. For example, the "volume" of the quadrangle of reference  $(-1 \leq \xi \leq +1, 1 < \eta < +1)$  is worth 4.

## 2 Linear elements: SE2, SE3 and SE4


**SE2** : segment with 2 nodes

many nodes : 2  
many nodes tops : 2

**SE3** : segment with 3 nodes

many nodes : 3  
many nodes tops : 2


	$x$
$N1$	-1.0
$N2$	1.0
$N3$	0.0



**SE4** : segment with 4 nodes

many nodes : 4  
many nodes tops : 2

	$x$
$N1$	-1.0
$N2$	1.0
$N3$	-1/3
$N4$	+1/3



functions of form of the segment with 2 nodes:

$$w_1(x) = 0.5(1-x) \quad w_2(x) = 0.5(1+x)$$

functions of form of the segment with 3 nodes:

$$w_1(x) = -0.5(1-x)x \quad w_2(x) = 0.5(1+x)x \quad w_3(x) = (1+x)(1-x)$$

functions of form of the segment with 4 nodes:

$$w_1(x) = \frac{9}{16}(1-x)\left(x+\frac{1}{3}\right)\left(x-\frac{1}{3}\right)$$

$$w_2(x) = -\frac{9}{16}(1+x)\left(\frac{1}{3}-x\right)\left(x+\frac{1}{3}\right)$$


$$w_3(x) = \frac{27}{16}(x-1)(x+1)\left(x-\frac{1}{3}\right)$$

$$w_4(x) = -\frac{27}{16}(x-1)(x+1)\left(x+\frac{1}{3}\right)$$

Many points of integration	Not	$x$	Weight
1	1	0.0	2.0
2	1	0.577350269189626	1.0
	2	-0.577350269189626	1.0
3	1	-0.774596669241	0.55555...
	2	0.0	0.88888...
	3	0.774596669241	0.55555...
4	1	0.339981043584856	0.652145154862546
	2	-0.339981043584856	0.652145154862546
	3	0.861136311594053	0.347854845137454
	4	-0.861136311594053	0.347854845137454

## 3 Surface elements

### 3.1 Triangles: ELREFE TR3, TR6, TR7



Coordinates of the nodes:

	$\xi$	$\eta$
N1	0.0	0.0
N2	1.0	0.0
N3	0.0	1.0
N4	0.5	0.0
N5	0.5	0.5
N6	0.0	0.5
N7	1/3	1/3

Family	Not	$\xi$	$\eta$	Weight
FPG1	1	1/3	1/3	1/2
FPG3	1	1/6	1/6	1/6
	2	2/3	1/6	1/6
	3	1/6	2/3	1/6
FPG4	1	1/5	1/5	25/(24*4)
	2	3/5	1/5	25/(24*4)
	3	1/5	3/5	25/(24*4)
	4	1/3	1/3	-27/(24*4)
FPG6	1	B	B	P2
	2	1 - 2b	B	P2
	3	B	1 - 2b	P2
	4	has	1 - 2a	P1
	5	has	has	P1
	6	1 - 2a	has	P1
COT3	1	1/2	1/2	1/6
	2	0	1/2	1/6
	3	1/2	0	1/6

With  $P1 = 0.11169079483905,$   $P2 = 0.0549758718227661,$   
 $With = 0.445948490915965,$   $B = 0.091576213509771$

Family	Not	$\xi$	$\eta$	Weight
FPG7	1	1/3	1/3	9/80
	2	With	With	P1
	3	1-2A	With	P1
	4	With	1-2A	P1
	5	B	B	P2
	6	1-2B	B	P2
	7	B	1-2B	P2

With  $With = 0.470142064105115$   
 $B = 0.101286507323456$   
 $P1 = 0.066197076394253$   
 $P2 = 0.062969590272413$

Family	Not	$\xi$	$\eta$	Weight
FPG12	1	With	With	P1
	2	1-2A	With	P1
	3	With	1-2A	P1
	4	B	B	P2
	5	1-2B	B	P2
	6	B	1-2B	P2
	7	C	D	P3
	8	D	C	P3
	9	1-C-D	C	P3
	10	1-C-D	D	P3
	11	C	1-C-D	P3
	12	D	1-C-D	P3

With                      With = 0.063089014491502  
 B = 0.249286745170910  
 C = 0.310352451033785  
 D = 0.053145049844816  
 P1 = 0.025422453185103  
 P2 = 0.058393137863189  
 P3 = 0.041425537809187

**TR3** : triangle with 3 nodes  
 many nodes                      : 3  
 many nodes tops                : 3

functions of form and derived first of the triangle with 3 nodes:

$N$	$\partial N / \partial \xi$	$\partial N / \partial \eta$
$1 - \xi - \eta$	-1	-1
$\xi$	1	0
$\eta$	0	1

**TR6** : triangle with 6 nodes  
 many nodes                      : 6  
 many nodes tops                : 3

functions of form, derived first of the triangle with 6 nodes:

$N$	$\partial N / \partial \xi$	$\partial N / \partial \eta$
$-(1 - \xi - \eta)(1 - 2(1 - \xi - \eta))$	$1 - 4(1 - \xi - \eta)$	$1 - 4(1 - \xi - \eta)$
$-\xi(1 - 2\xi)$	$-1 + 4\xi$	0
$-\eta(1 - 2\eta)$	0	$-1 + 4\eta$
$4\xi(1 - \xi - \eta)$	$4(1 - 2\xi - \eta)$	$-4\xi$
$4\xi\eta$	$4\eta$	$4\xi$
$4\eta(1 - \xi - \eta)$	$-4\eta$	$4(1 - \xi - 2\eta)$



derived seconds from the triangle with 6 nodes:

$\partial^2 N / \partial \xi^2$	$\partial^2 N / \partial \xi \partial \eta$	$\partial^2 N / \partial \eta^2$
4	4	4
4	0	0
0	0	4
-8	-4	0
0	4	0
0	-4	-8

**TR7** : triangle with 7 nodes

many nodes : 7  
many nodes tops : 3

functions of form of the triangle with 7 nodes:

$N$
$1 - 3(\xi + \eta) + 2(\xi^2 + \eta^2) + 7\xi\eta - 3\xi\eta(\xi + \eta)$
$\xi(-1 + 2\xi + 3\eta - 3\eta(\xi + \eta))$
$\eta(-1 + 2\xi + 3\eta - 3\xi(\xi + \eta))$
$4\xi(1 - \xi - 4\eta + 3\eta(\xi + \eta))$
$4\xi\eta(-2 + 3(\xi + \eta))$
$4\eta(1 - 4\xi - \eta + 3\xi(\xi + \eta))$
$27\xi\eta(1 - \xi - \eta)$


derived first from the triangle with 7 nodes:

$\partial N / \partial \xi$	$\partial N / \partial \eta$
$-3 + 4\xi + 7\eta - 6\xi\eta - 3\eta^2$	$-3 + 7\xi + 4\eta - 6\xi\eta - 3\xi^2$
$-1 + 4\xi + 3\eta - 6\xi\eta - 3\eta^2$	$3\xi(1 - \xi - 2\eta)$
$3\xi(1 - 2\eta - \xi)$	$-1 + 3\xi + 4\eta - 6\xi\eta - 3\xi^2$
$4(1 - 2\xi - 4\eta + 6\xi\eta + 3\eta^2)$	$4\xi(-4 + 3\xi + 6\eta)$
$4\eta(-2 + 6\xi + 3\eta)$	$4\xi(-2 + 3\xi + 6\eta)$
$4\eta(-4 + 6\xi + 3\eta)$	$4(-1 - 4\xi - 2\eta + 6\xi\eta + 3\xi^2)$
$27\eta(1 - 2\xi - \eta)$	$27\xi(1 - \xi - 2\eta)$

derived seconds from the triangle with 7 nodes:

$\partial^2 N / \partial \xi^2$	$\partial^2 N / \partial \xi \partial \eta$	$\partial^2 N / \partial \eta^2$
$4 - 6\eta$	$7 - 6\xi - 6\eta$	$4 - 6\xi$
$4 - 6\eta$	$3 - 6\xi - 6\eta$	$-6\xi$
$-6\eta$	$3 - 6\xi - 6\eta$	$4 - 6\xi$
$4(-2 + 6\eta)$	$4(-4 + 6\xi + 6\eta)$	$24\xi$
$24\eta$	$4(-2 + 6\xi + 6\eta)$	$24\xi$
$24\eta$	$4(-4 + 6\xi + 6\eta)$	$4(-2 + 6\xi)$
$-54\eta$	$27(1 - 2\xi - 2\eta)$	$-54\xi$

## 3.2 Quadrangles: ELREFE QU4, QU8, QU9



Coordinates of the nodes:

	$\xi$	$\eta$
N1	-1.0	-1.0
N2	1.0	-1.0
N3	1.0	1.0
N4	-1.0	1.0
N5	0.0	-1.0
N6	1.0	0.0
N7	0.0	1.0
N8	-1.0	0.0
N9	0.0	0.0

Family	Not	$\xi$	$\eta$	Weight
FPG1	1	0	0	4
FPG4	1	$-a$	$-a$	1.0
	2	$a$	$-a$	1.0
	3	$a$	$a$	1.0
	4	$-a$	$a$	1.0
		$a=1/\sqrt{3}$		
FPG9	1	$-a$	$-a$	25/81
	2	$a$	$-a$	25/81
	3	$a$	$a$	25/81
	4	$-a$	$a$	25/81
	5	0.0	$-a$	40/81
	6	$a$	0.0	40/81
	7	0.0	has	40/81
	8	$-a$	0.0	40/81
	9	0.0	0.0	64/81
		$a=0.774596669241483$		

**QU4** : quadrangle with 4 nodes

many nodes : 4  
many nodes tops : 4

functions of form, derived first and seconds of the quadrangle with 4 nodes:

$N$	$\partial N / \partial \xi$	$\partial N / \partial \eta$
$(1-\xi)(1-\eta)/4$	$-(1-\eta)/4$	$-(1-\xi)/4$
$(1+\xi)(1-\eta)/4$	$(1-\eta)/4$	$-(1+\xi)/4$
$(1+\xi)(1+\eta)/4$	$(1+\eta)/4$	$(1+\xi)/4$
$(1-\xi)(1+\eta)/4$	$-(1+\eta)/4$	$(1-\xi)/4$

$\partial^2 N / \partial \xi^2$	$\partial^2 N / \partial \xi \partial \eta$	$\partial^2 N / \partial \eta^2$
0	1/4	0
0	-1/4	0
0	1/4	0
0	-1/4	0

**QU8** : quadrangle with 8 nodes

many nodes : 8  
many nodes tops : 4

functions of form and derived first of the quadrangle with 8 nodes:

$N$	$\partial N / \partial \xi$	$\partial N / \partial \eta$
$(1-\xi)(1-\eta)(-1-\xi-\eta)/4$	$(1-\eta)(2\xi+\eta)/4$	$(1-\xi)(\xi+2\eta)/4$
$(1+\xi)(1-\eta)(-1+\xi-\eta)/4$	$(1-\eta)(2\xi-\eta)/4$	$-(1+\xi)(\xi-2\eta)/4$
$(1+\xi)(1+\eta)(-1+\xi+\eta)/4$	$(1+\eta)(2\xi+\eta)/4$	$(1+\xi)(\xi+2\eta)/4$
$(1-\xi)(1+\eta)(-1-\xi+\eta)/4$	$-(1+\eta)(-2\xi+\eta)/4$	$(1-\xi)(-\xi+2\eta)/4$
$(1-\xi^2)(1-\eta)/2$	$-\xi(1-\eta)$	$-(1-\xi^2)/2$
$(1+\xi)(1-\eta^2)/2$	$(1-\eta^2)/2$	$-\eta(1+\xi)$
$(1-\xi^2)(1+\eta)/2$	$-\xi(1+\eta)$	$(1-\xi^2)/2$
$(1-\xi)(1-\eta^2)/2$	$-(1-\eta^2)/2$	$-\eta(1-\xi)$

derived seconds from the quadrangle with 8 nodes:

$\partial^2 N / \partial \xi^2$	$\partial^2 N / \partial \xi \partial \eta$	$\partial^2 N / \partial \eta^2$
$(1-\eta)/2$	$(1-2\xi-2\eta)/4$	$(1-\xi)/2$
$(1-\eta)/2$	$-(1+2\xi-2\eta)/4$	$(1+\xi)/2$
$(1+\eta)/2$	$(1+2\xi+2\eta)/4$	$(1+\xi)/2$
$(1+\eta)/2$	$-(1-2\xi+2\eta)/4$	$(1-\xi)/2$
$-1+\eta$	$\xi$	0
0	$-\eta$	$-1-\xi$
$-1-\eta$	$-\xi$	0
0	$\eta$	$-1+\xi$

**QU9** : quadrangle with 9 nodes

many nodes : 9  
many nodes tops : 4

functions of form and derived first of the quadrangle with 9 nodes:


$[N]$	$[\partial N/\partial \xi]$	$[\partial N/\partial \eta]$
$\xi\eta(\xi-1)(\eta-1)/4$	$(2\xi-1)\eta(\eta-1)/4$	$\xi(\xi-1)(2\eta-1)/4$
$\xi\eta(\xi+1)(\eta-1)/4$	$(2\xi+1)\eta(\eta-1)/4$	$\xi(\xi+1)(2\eta-1)/4$
$\xi\eta(\xi+1)(\eta+1)/4$	$(2\xi+1)\eta(\eta+1)/4$	$\xi(\xi+1)(2\eta+1)/4$
$\xi\eta(\xi-1)(\eta+1)/4$	$(2\xi-1)\eta(\eta+1)/4$	$\xi(\xi-1)(2\eta+1)/4$
$(1-\xi^2)\eta(\eta-1)/2$	$-\xi\eta(\eta-1)$	$(1-\xi^2)(2\eta-1)/2$
$\xi(\xi+1)(1-\eta^2)/2$	$(2\xi+1)(1-\eta^2)/2$	$-\xi\eta(\xi+1)$
$(1-\xi^2)\eta(\eta+1)/2$	$-\xi\eta(\eta+1)$	$(1-\xi^2)(2\eta+1)/2$
$\xi(\xi-1)(1-\eta^2)/2$	$(2\xi-1)(1-\eta^2)/2$	$-\xi\eta(\xi-1)$
$(1-\xi^2)(1-\eta^2)$	$-2\xi(1-\eta^2)$	$-2\eta(1-\xi^2)$

derived seconds from the quadrangle with 9 nodes:

$[\partial^2 N/\partial \xi^2]$	$[\partial^2 N/\partial \xi \partial \eta]$	$[\partial^2 N/\partial \eta^2]$
$\eta(\eta-1)/2$	$(\xi-1/2)(\eta-1/2)$	$\xi(\xi-1)/2$
$\eta(\eta-1)/2$	$(\xi+1/2)(\eta-1/2)$	$\xi(\xi+1)/2$
$\eta(\eta+1)/2$	$(\xi+1/2)(\eta+1/2)$	$\xi(\xi+1)/2$
$\eta(\eta+1)/2$	$(\xi-1/2)(\eta+1/2)$	$\xi(\xi-1)/2$
$-\eta(\eta-1)$	$-\xi(2\eta-1)$	$1-\xi^2$
$1-\eta^2$	$-\eta(2\xi+1)$	$-\xi(\xi+1)$
$-\eta(\eta+1)$	$-\xi(2\eta+1)$	$1-\xi^2$
$1-\eta^2$	$-\eta(2\xi-1)$	$-\xi(\xi-1)$
$-2(1-\eta^2)$	$4\xi\eta$	$-2(1-\xi^2)$

## 4 Voluminal elements

### 4.1 Tetrahedrons: ELREFE TE4, T10



Coordinates of the nodes:

	$x$	$y$	$z$
$N1$	0.	1.	0.
$N2$	0.	0.	1.
$N3$	0.	0.	0.
$N4$	1.	0.	0.
$N5$	0.	0.5	0.5
$N6$	0.	0.	0.5
$N7$	0.	0.5	0.
$N8$	0.5	0.5	0.
$N9$	0.5	0.	0.5
$N10$	0.5	0.	0.

**Functions of form:**

Formula with 4 nodes

$$\begin{cases} w_1(x, y, z) = y \\ w_2(x, y, z) = z \\ w_3(x, y, z) = 1 - x - y - z \\ w_4(x, y, z) = x \end{cases}$$

Formula with 10 nodes

$$w_1 = y(2y - 1)$$

$$w_2 = z(2z - 1)$$

$$w_3 = (1 - x - y - z)(1 - 2x - 2y - 2z)$$

$$w_4 = x(2x - 1)$$

$$w_5 = 4yz$$

$$w_6 = 4z(1 - x - y - z)$$

$$w_7 = 4y(1 - x - y - z)$$

$$w_8 = 4xy$$

$$w_9 = 4xz$$

$$w_{10} = 4x(1 - x - y - z)$$

**Digital formula of integration:**

Formula with 1 point, of order 1 in  $x, y, z$  : (FPG1)

Not	$x$	$y$	$z$	Weight
1	1/4	1/4	1/4	1/6

Formula at 4 points, of order 2 in  $x, y, z$  : (FPG4)

Not	$x$	$y$	$z$	Weight
1	$a$	$a$	$a$	1/24
2	$a$	$a$	$b$	1/24
3	$a$	$b$	$a$	1/24
4	$b$	$a$	$a$	1/24

with:  $a = \frac{5 - \sqrt{5}}{20}$ ,  $b = \frac{5 + 3\sqrt{5}}{20}$

Formula at 5 points, of order 3 in  $x, y, z$  : (FPG5)

Not	$x$	$y$	$z$	Weight
1	$a$	$a$	$a$	-2/15
2	$b$	$b$	$b$	3/40
3	$b$	$b$	$c$	3/40
4	$b$	$c$	$b$	3/40
5	$c$	$b$	$b$	3/40

With:  $a = 0.25$ ,  $b = \frac{1}{6}$ ,  $c = 0.5$

Formula at 15 points, of order 5 in  $x, y, z$  : (FPG15)

Not	$x$	$y$	$z$	Weight
1	$a$	$a$	$a$	8/405
2	$b_1$	$b_1$	$b_1$	$\frac{2665-14\sqrt{15}}{226800}$
3	$b_1$	$b_1$	$c_1$	
4	$b_1$	$c_1$	$b_1$	
5	$c_1$	$b_1$	$b_1$	
6	$b_2$	$b_2$	$b_2$	
7	$b_2$	$b_2$	$c_2$	$\frac{2665+14\sqrt{15}}{226800}$
8	$b_2$	$c_2$	$b_2$	
9	$c_2$	$b_2$	$b_2$	
10	$d$	$d$	$e$	$\frac{5}{567}$
11	$d$	$e$	$d$	
12	$e$	$d$	$d$	
13	$d$	$e$	$e$	
14	$e$	$d$	$e$	
15	$e$	$e$	$d$	

with:  
 $a=0.25$

$$b_1 = \frac{7+\sqrt{15}}{34}$$

$$b_2 = \frac{7-\sqrt{15}}{34}$$


$$c_1 = \frac{13-3\sqrt{15}}{34}$$

$$c_2 = \frac{13+3\sqrt{15}}{34}$$

$$d = \frac{5-\sqrt{15}}{20}$$

$$e = \frac{5+\sqrt{15}}{20}$$

## 4.2 Pentahedrons: ELREFE PE6, P15, P18



Coordinates of the nodes:

	$x$	$y$	$z$
$N1$	-1.	1.	0.
$N2$	-1.	0.	1.
$N3$	-1.	0.	0.
$N4$	1.	1.	0.
$N5$	1.	0.	1.
$N6$	1.	0.	0.
$N7$	-1.	0.5	0.5.
$N8$	-1.	0.	0.5.
$N9$	-1.	0.5	0.
$N10$	0.	1.	0.
$N11$	0.	0.	1.
$N12$	0.	0.	0.
$N13$	1.	0.5	0.5
$N14$	1.	0.	0.5
$N15$	1.	0.5	0.
$N16$	0.	0.5	0.5
$N17$	0.	0.	0.5
$N18$	0.	0.5	0.

Functions of form:

Formula with 6 nodes

$$w_1 = \frac{1}{2} y (1-x)$$

$$w_4 = \frac{1}{2} y (x+1)$$

$$w_2 = \frac{1}{2} z (1-x)$$

$$w_5 = \frac{1}{2} z (x+1)$$

$$w_3 = \frac{1}{2} (1-y-z)(1-x)$$

$$w_6 = \frac{1}{2} (1-y-z)(x+1)$$

Formula with 15 nodes

$$w_1 = y(1-x)(2y-2-x)/2$$

$$w_9 = 2y(1-y-z)(1-x)$$

$$w_2 = z(1-x)(2z-2-x)/2$$

$$w_{10} = y(1-x^2)$$

$$w_3 = (x-1)(1-y-z)(x+2y+2z)/2$$

$$w_{11} = z(1-x^2)$$

$$w_4 = y(1+x)(2y-2+x)/2$$

$$w_{12} = (1-y-z)(1-x^2)$$

$$w_5 = z(1+x)(2z-2+x)/2$$

$$w_{13} = 2yz(1+x)$$

$$w_6 = (-x-1)(1-y-z)(-x+2y+2z)/2$$

$$w_{14} = 2z(1-y-z)(1+x)$$

$$w_7 = 2yz(1-x)$$

$$w_{15} = 2y(1-y-z)(1+x)$$

$$w_8 = 2z(1-y-z)(1-x)$$



Formula with 18 nodes

$$\begin{aligned}
 w_1 &= x y (x-1)(2y-1)/2 & w_{10} &= y(1-x^2)(2y-1) \\
 w_2 &= x z (x-1)(2z-1)/2 & w_{11} &= z(1-x^2)(2z-1) \\
 w_3 &= x(x-1)(z+y-1)(2z+2y-1)/2 & w_{12} &= (1-x^2)(z+y-1)(2z+2y-1) \\
 w_4 &= x y (x+1)(2y-1)/2 & w_{13} &= 2 x y z (x+1) \\
 w_5 &= x z (x+1)(2z-1)/2 & w_{14} &= -2 x z (x+1)(z+y-1) \\
 w_6 &= x(x+1)(z+y-1)(2z+2y-1)/2 & w_{15} &= -2 x y (x+1)(z+y-1) \\
 w_7 &= 2 x y z (x-1) & w_{16} &= 4 y z (1-x^2) \\
 w_8 &= -2 x z (x-1)(z+y-1) & w_{17} &= 4 z (x^2-1)(z+y-1) \\
 w_9 &= -2 x y (x-1)(z+y-1) & w_{18} &= 4 y (x^2-1)(z+y-1)
 \end{aligned}$$

**Formulas of digital integration at 6 points** (order 3 in  $x$ , order 2 in  $y$  and  $z$ ) (FPG6)

Not	$x$	$y$	$z$	Weight
1	$-1/\sqrt{3}$	0.5	0.5	1/6
2	$-1/\sqrt{3}$	0.	0.5	1/6
3	$-1/\sqrt{3}$	0.5	0.	1/6
4	$1/\sqrt{3}$	0.5	0.5	1/6
5	$1/\sqrt{3}$	0.	0.5	1/6
6	$1/\sqrt{3}$	0.5	0.	1/6

**Digital formula of integration at 8 points: (FPG8)**

2 points of Gauss in  $x$  (order 3).

4 points of Hammer in  $y$  and  $z$  (order 3).

Not	$x$	$y$	$z$	Weight
1	$-a$	1/3	1/3	$-27/96$
2	$-a$	0.6	0.2	$25/96$
3	$-a$	0.2	0.6	$25/96$
4	$-a$	0.2	0.2	$25/96$
5	$+a$	1/3	1/3	$-27/96$
6	$+a$	0.6	0.2	$25/96$
7	$+a$	0.2	0.6	$25/96$
8	$+a$	0.2	0.2	$25/96$

With  $a=0.577350269189626$

## Digital formula of integration at 21 points: (FPG21)

3 points of Gauss in  $x$  (order 5).


7 points of Hammer in  $y$  and  $z$  (order 5 in  $y$  and  $z$ ).

Not	$x$	$y$	$z$	Weight
1	$-\alpha$	$1/3$	$1/3$	$c_1 \frac{9}{80}$
2	$-\alpha$	$a$	$a$	$c_1 \left( \frac{155 + \sqrt{15}}{2400} \right)$
3	$-\alpha$	$1-2a$	$a$	
4	$-\alpha$	$a$	$1-2a$	
5	$-\alpha$	$b$	$b$	$c_1 \left( \frac{155 - \sqrt{15}}{2400} \right)$
6	$-\alpha$	$1-2b$	$b$	
7	$-\alpha$	$b$	$1-2b$	
8	0.	$1/3$	$1/3$	$c_2 \frac{9}{80}$
9	0.	$a$	$a$	$c_2 \left( \frac{155 + \sqrt{15}}{2400} \right)$
10	0.	$1-2a$	$a$	
11	0.	$a$	$1-2a$	
12	0.	$b$	$b$	$c_2 \left( \frac{155 - \sqrt{15}}{2400} \right)$
13	0.	$1-2b$	$b$	
14	0.	$b$	$1-2b$	
15	$\alpha$	$1/3$	$1/3$	$c_1 \frac{9}{80}$
16	$\alpha$	$b$	$a$	$c_1 \left( \frac{155 + \sqrt{15}}{2400} \right)$
17	$\alpha$	$1-2a$	$a$	
18	$\alpha$	$a$	$1-2a$	
19	$\alpha$	$b$	$b$	$c_1 \left( \frac{155 - \sqrt{15}}{2400} \right)$
20	$\alpha$	$1-2b$	$b$	
21	$\alpha$	$b$	$1-2b$	

with:

$$\alpha = \sqrt{\frac{3}{5}} \quad c_1 = \frac{5}{9} \quad c_2 = \frac{8}{9} \quad a = \frac{6 + \sqrt{15}}{21} \quad b = \frac{6 - \sqrt{15}}{21}$$

## 4.3 Hexahedrons: ELREFE HE8 , H20 , H27



Coordinates of the nodes:

	x	y	z
N1	-1.	-1.	-1.
N2	1.	-1.	-1.
N3	1.	1.	-1.
N4	-1.	1.	-1.
N5	-1.	-1.	1.
N6	1.	-1.	1.
N7	1.	1.	1.
N8	-1.	1.	1.
N9	0.	-1.	-1.
N10	1.	0.	-1.
N11	0.	1.	-1.
N12	-1.	0.	-1.
N13	-1.	-1.	0.
N14	1.	-1.	0.
N15	1.	1.	0.
N16	-1.	1.	0.
N17	0.	-1.	1.
N18	1.	0.	1.
N19	0.	1.	1.
N20	-1.	0.	1.
N21	0.	0.	-1.
N22	0.	-1.	0.
N23	1.	0.	0.
N24	0.	1.	0.
N25	-1.	0.	0.
N26	0.	0.	1.
N27	0.	0.	0.

Functions of form:

Formula with 8 nodes

$$w_1 = \frac{1}{8}(1-x)(1-y)(1-z)$$

$$w_5 = \frac{1}{8}(1-x)(1-y)(1+z)$$

$$w_2 = \frac{1}{8}(1+x)(1-y)(1-z)$$

$$w_6 = \frac{1}{8}(1+x)(1-y)(1+z)$$

$$w_3 = \frac{1}{8}(1+x)(1+y)(1-z)$$

$$w_7 = \frac{1}{8}(1+x)(1+y)(1+z)$$

$$w_4 = \frac{1}{8}(1-x)(1+y)(1-z)$$

$$w_8 = \frac{1}{8}(1-x)(1+y)(1+z)$$

Formula with 20 nodes

$$w_1 = \frac{1}{8}(1-x)(1-y)(1-z)(-2-x-y-z)$$

$$w_2 = \frac{1}{8}(1+x)(1-y)(1-z)(-2+x-y-z)$$

$$w_3 = \frac{1}{8}(1+x)(1+y)(1-z)(-2+x+y-z)$$

$$w_4 = \frac{1}{8}(1-x)(1+y)(1-z)(-2-x+y-z)$$

$$w_5 = \frac{1}{8}(1-x)(1-y)(1+z)(-2-x-y+z)$$

$$w_6 = \frac{1}{8}(1+x)(1-y)(1+z)(-2+x-y+z)$$

$$w_7 = \frac{1}{8}(1+x)(1+y)(1+z)(-2+x+y+z)$$

$$w_8 = \frac{1}{8}(1-x)(1+y)(1+z)(-2-x+y+z)$$

$$w_9 = \frac{1}{4}(1-x^2)(1-y)(1-z)$$

$$w_{10} = \frac{1}{4}(1-y^2)(1+x)(1-z)$$

$$w_{11} = \frac{1}{4}(1-x^2)(1+y)(1-z)$$

$$w_{12} = \frac{1}{4}(1-y^2)(1-x)(1-z)$$

$$w_{13} = \frac{1}{4}(1-z^2)(1-x)(1-y)$$

$$w_{14} = \frac{1}{4}(1-z^2)(1+x)(1-y)$$

$$w_{15} = \frac{1}{4}(1-z^2)(1+x)(1+y)$$

$$w_{16} = \frac{1}{4}(1-z^2)(1-x)(1+y)$$

$$w_{17} = \frac{1}{4}(1-x^2)(1-y)(1+z)$$

$$w_{18} = \frac{1}{4}(1-y^2)(1+x)(1+z)$$

$$w_{19} = \frac{1}{4}(1-x^2)(1+y)(1+z)$$

$$w_{20} = \frac{1}{4}(1-y^2)(1-x)(1+z)$$

Formula with 27 nodes

$$\begin{aligned}
 w_1 &= \frac{1}{8} x(x-1) y(y-1) z(z-1) & w_{15} &= \frac{1}{4} x(x+1) y(y+1) (1-z^2) \\
 w_2 &= \frac{1}{8} x(x+1) y(y-1) z(z-1) & w_{16} &= \frac{1}{4} x(x-1) y(y+1) (1-z^2) \\
 w_3 &= \frac{1}{8} x(x+1) y(y+1) z(z-1) & w_{17} &= \frac{1}{4} (1-x^2) y(y-1) z(z+1) \\
 w_4 &= \frac{1}{8} x(x-1) y(y+1) z(z-1) & w_{18} &= \frac{1}{4} x(x+1) (1-y^2) z(z+1) \\
 w_5 &= \frac{1}{8} x(x-1) y(y-1) z(z+1) & w_{19} &= \frac{1}{4} (1-x^2) y(y+1) z(z+1) \\
 w_6 &= \frac{1}{8} x(x+1) y(y-1) z(z+1) & w_{20} &= \frac{1}{4} x(x-1) (1-y^2) z(z+1) \\
 w_7 &= \frac{1}{8} x(x+1) y(y+1) z(z+1) & w_{21} &= \frac{1}{2} (1-x^2) (1-y^2) z(z-1) \\
 w_8 &= \frac{1}{8} x(x-1) y(y+1) z(z+1) & w_{22} &= \frac{1}{2} (1-x^2) y(y-1) (1-z^2) \\
 w_9 &= \frac{1}{4} (1-x^2) y(y-1) z(z-1) & w_{23} &= \frac{1}{2} x(x+1) (1-y^2) (1-z^2) \\
 w_{10} &= \frac{1}{4} x(x+1) (1-y^2) z(z-1) & w_{24} &= \frac{1}{2} (1-x^2) y(y+1) (1-z^2) \\
 w_{11} &= \frac{1}{4} (1-x^2) y(y+1) z(z-1) & w_{25} &= \frac{1}{2} x(x-1) (1-y^2) (1-z^2) \\
 w_{12} &= \frac{1}{4} x(x-1) (1-y^2) z(z-1) & w_{26} &= \frac{1}{2} (1-x^2) (1-y^2) z(z+1) \\
 w_{13} &= \frac{1}{4} x(x-1) y(y-1) (1-z^2) & w_{27} &= (1-x^2) (1-y^2) (1-z^2) \\
 w_{14} &= \frac{1}{4} x(x+1) y(y-1) (1-z^2)
 \end{aligned}$$

Formula of squaring of Gauss at 2 points in each direction (order 3) (FPG8)

Not	x	y	z	Weight
1	$-1/\sqrt{3}$	$-1/\sqrt{3}$	$-1/\sqrt{3}$	1.
2	$-1/\sqrt{3}$	$-1/\sqrt{3}$	$1/\sqrt{3}$	1.
3	$-1/\sqrt{3}$	$1/\sqrt{3}$	$-1/\sqrt{3}$	1.
4	$-1/\sqrt{3}$	$1/\sqrt{3}$	$1/\sqrt{3}$	1.
5	$1/\sqrt{3}$	$-1/\sqrt{3}$	$-1/\sqrt{3}$	1.
6	$1/\sqrt{3}$	$-1/\sqrt{3}$	$1/\sqrt{3}$	1.
7	$1/\sqrt{3}$	$1/\sqrt{3}$	$-1/\sqrt{3}$	1.
8	$1/\sqrt{3}$	$1/\sqrt{3}$	$1/\sqrt{3}$	1.

**Formula of squaring of Gauss at 3 points in each direction (order 5): ( FPG27 )**

Not	x	y	z	Weight
1	$-\alpha$	$-\alpha$	$-\alpha$	$c_1^3$
2	$-\alpha$	$-\alpha$	0.	$c_1^2 c_2$
3	$-\alpha$	$-\alpha$	$\alpha$	$c_1^3$
4	$-\alpha$	0.	$-\alpha$	$c_1^2 c_2$
5	$-\alpha$	0.	0.	$c_1 c_2^2$
6	$-\alpha$	0.	$\alpha$	$c_1^2 c_2$
7	$-\alpha$	$\alpha$	$-\alpha$	$c_1^3$
8	$-\alpha$	$\alpha$	0.	$c_1^2 c_2$
9	$-\alpha$	$\alpha$	$\alpha$	$c_1^3$
10	0.	$-\alpha$	$-\alpha$	$c_1^2 c_2$
11	0.	$-\alpha$	0.	$c_1 c_2^2$
12	0.	$-\alpha$	$\alpha$	$c_1^2 c_2$
13	0.	0.	$-\alpha$	$c_1 c_2^2$
14	0.	0.	0.	$c_2^3$
15	0.	0.	$\alpha$	$c_1 c_2^2$
16	0.	$\alpha$	$-\alpha$	$c_1^2 c_2$
17	0.	$\alpha$	0.	$c_1 c_2^2$
18	0.	$\alpha$	$\alpha$	$c_1^2 c_2$
19	$\alpha$	$-\alpha$	$-\alpha$	$c_1^3$
20	$\alpha$	$-\alpha$	0.	$c_1^2 c_2$
21	$\alpha$	$-\alpha$	$\alpha$	$c_1^3$
22	$\alpha$	0.	$-\alpha$	$c_1^2 c_2$
23	$\alpha$	0.	0.	$c_1 c_2^2$
24	$\alpha$	0.	$\alpha$	$c_1^2 c_2$
25	$\alpha$	$\alpha$	$-\alpha$	$c_1^3$
26	$\alpha$	$\alpha$	0.	$c_1^2 c_2$
27	$\alpha$	$\alpha$	$\alpha$	$c_1^3$


with:

$$\alpha = \sqrt{\frac{3}{5}}$$

$$c_1 = \frac{5}{9}$$

$$c_2 = \frac{8}{9}$$

## 4.4 Pyramids: ELREFE PY5, P13



The square base is consisted the quadrangle  $N_1 N_2 N_3 N_4$  and  $N_5$  is the top of the pyramid.

	$x$	$y$	$z$
$N_1$	1.	0.	0.
$N_2$	0.	1.	0.
$N_3$	-1.	0.	0.
$N_4$	0.	-1.	0.
$N_5$	0.	0.	1.
$N_6$	0.5	0.5	0.
$N_7$	-0.5	0.5	0.
$N_8$	-0.5	-0.5	0.
$N_9$	0.5	-0.5	0.
$N_{10}$	0.5	0.	0.5
$N_{11}$	0.	0.5	0.5
$N_{12}$	-0.5	0.	0.5
$N_{13}$	0.	-0.5	0.5



## Functions of form:

### Formula with 5 nodes

$$w_1 = \frac{(-x+y+z-1)(-x-y+z-1)}{4(1-z)}$$

$$w_2 = \frac{(-x-y+z-1)(x-y+z-1)}{4(1-z)}$$

$$w_3 = \frac{(x+y+z-1)(x-y+z-1)}{4(1-z)}$$

$$w_4 = \frac{(x+y+z-1)(-x+y+z-1)}{4(1-z)}$$

$$w_5 = z$$

### Formula with 13 nodes

$$w_1 = \frac{(-x+y+z-1)(-x-y+z-1)(x-0.5)}{2(1-z)}$$

$$w_2 = \frac{(-x-y+z-1)(x-y+z-1)(y-0.5)}{2(1-z)}$$

$$w_3 = \frac{(x-y+z-1)(x+y+z-1)(-x-0.5)}{2(1-z)}$$

$$w_4 = \frac{(x+y+z-1)(-x+y+z-1)(-y-0.5)}{2(1-z)}$$

$$w_5 = 2z(z-0.5)$$

$$w_6 = -\frac{(-x+y+z-1)(-x-y+z-1)(x-y+z-1)}{2(1-z)}$$

$$w_7 = -\frac{(-x-y+z-1)(x-y+z-1)(x+y+z-1)}{2(1-z)}$$

$$w_8 = -\frac{(x-y+z-1)(x+y+z-1)(-x+y+z-1)}{2(1-z)}$$

$$w_9 = -\frac{(x+y+z-1)(-x+y+z-1)(-x-y+z-1)}{2(1-z)}$$

$$w_{10} = \frac{z(-x+y+z-1)(-x-y+z-1)}{1-z}$$

$$w_{11} = \frac{z(-x-y+z-1)(x-y+z-1)}{1-z}$$

$$w_{12} = \frac{z(x-y+z-1)(x+y+z-1)}{1-z}$$

$$w_{13} = \frac{z(x+y+z-1)(-x+y+z-1)}{1-z}$$

Digital formula of integration at 5 points (FPG5):

Not	x	y	z	Weight
1	0.5	0.	H <sub>1</sub>	2/15
2	0.	0.5	H <sub>1</sub>	2/15
3	-0.5	0.	H <sub>1</sub>	2/15
4	0.	-0.5	H <sub>1</sub>	2/15
5	0.	0.	H <sub>2</sub>	2/15

with:

$$h_1 = 0.1531754163448146$$

$$h_2 = 0.6372983346207416$$

Digital formula of integration at 6 points (FPG6):

Not	x	y	z	Weight
1	has	0.	$h_1$	$p_1$
2	0.	$a$	$h_1$	$p_1$
3	$-a$	0.	$h_1$	$p_1$
4	0.	$-a$	$h_1$	$p_1$
5	0.	0.	$h_2$	$p_2$
6	0.	0.	$h_3$	$p_3$

with:

$$p_1 = 0.1024890634400000$$

$$p_2 = 0.1100000000000000$$

$$p_3 = 0.1467104129066667$$

$$a = 0.5702963741068025$$

$$h_1 = 0.1666666666666666$$

$$h_2 = 0.08063183038464675$$

$$h_3 = 0.6098484849057127$$

Digital formula of integration at 27 points (FPG27):

Not	x	y	z	Weight
1	0.	0.	1/2	$a_1$
2	$b_1(1-z)/2$	$b_1(1-z)/2$	1/2	$b_6$
3	$-b_1(1-z)/2$	$b_1(1-z)/2$	1/2	$b_6$
4	$-b_1(1-z)/2$	$-b_1(1-z)/2$	1/2	$b_6$
5	$b_1(1-z)/2$	$-b_1(1-z)/2$	1/2	$b_6$
6	0.	0.	$1-b_1/2$	$b_6$
7	0.	0.	$1+b_1/2$	$b_6$
8	$c_1(1-z)$	0.	$(1-c_1)/2$	$c_8$
9	0.	$c_1(1-z)$	$(1-c_1)/2$	$c_8$
10	$-c_1(1-z)$	0.	$(1-c_1)/2$	$c_8$
11	0.	$-c_1(1-z)$	$(1-c_1)/2$	$c_8$
12	$c_1(1-z)$	0.	$(1+c_1)/2$	$c_8$
13	0.	$c_1(1-z)$	$(1+c_1)/2$	$c_8$
14	$-c_1(1-z)$	0.	$(1+c_1)/2$	$c_8$
15	0.	$-c_1(1-z)$	$(1+c_1)/2$	$c_8$
16	$d_1(1-z)/2$	$d_1(1-z)/2$	$(1-d_1)/2$	$d_{12}$
17	$-d_1(1-z)/2$	$d_1(1-z)/2$	$(1-d_1)/2$	$d_{12}$
18	$-d_1(1-z)/2$	$-d_1(1-z)/2$	$(1-d_1)/2$	$d_{12}$
19	$d_1(1-z)/2$	$-d_1(1-z)/2$	$(1-d_1)/2$	$d_{12}$
20	$d_1(1-z)$	0.	1/2	$d_{12}$
21	0.	$d_1(1-z)$	1/2	$d_{12}$
22	$-d_1(1-z)$	0.	1/2	$d_{12}$
23	0.	$-d_1(1-z)$	1/2	$d_{12}$
24	$d_1(1-z)/2$	$d_1(1-z)/2$	$(1+d_1)/2$	$d_{12}$
25	$-d_1(1-z)/2$	$d_1(1-z)/2$	$(1+d_1)/2$	$d_{12}$
26	$-d_1(1-z)/2$	$-d_1(1-z)/2$	$(1+d_1)/2$	$d_{12}$
27	$d_1(1-z)/2$	$-d_1(1-z)/2$	$(1+d_1)/2$	$d_{12}$

with:

$$\begin{aligned}
 a_1 &= 0.788073483 \\
 b_6 &= 0.499369002 \\
 b_1 &= 0.848418011 \\
 c_8 &= 0.478508449 \\
 c_1 &= 0.652816472 \\
 d_{12} &= 0.032303742 \\
 d_1 &= 1.106412899
 \end{aligned}$$

## 5 Bibliography

---

- 1 DHATT G., TOUZOT G.: A presentation of the finite element method 2<sup>ème</sup> edition. Editor: MALOINE S.A. Year 984

## 6 History of the versions of the document

---

Index Doc.	Version Aster	Author (S) or contributor (S), organization	Description of the modifications
E	8.4	J.Pellet, X.Desroches, EDF/R & D	Version 8 complete.
F	9.2	J.Pellet EDF/R & D /AMA	Correction concerning the HEXA27, cf drives REX 11036
F	9.4	J.Pellet EDF/R & D /AMA	Correction page 21 of the function of form w5 of the HEXA27 (card 12170)
			Correction function of form SE4 (card 26110)